BEAM THEORY – GEOMETRICAL PROPERTIES OF THE CROSS SECTION

STRUCTURAL MECHANICS

The ERAMCA Project

Environmental Risk Assessment and Mitigation on Cultural Heritage assets in Central Asia

V2022317

This work is licensed under a Creative Commons "Attribution-ShareAlike 4.0 International" license.







Lecturer/students objectives

Introduction

Centroid, first and second moments of area

Parallel-axis theorems

Principal axes

Simple sections

Composite areas

Examples





LECTURER/STUDENTS OBJECTIVES





- Present the mathematical instrument necessary to evaluate the influence of the cross section on the stress and deformation of beams.
- Learn the procedures necessary to the calculation of the geometrical properties of plane areas (centroid, first and second moments, principal axes). Apply the procedure to plane areas composed by elementary figures.





INTRODUCTION





The geometry of plane areas is examined to derive some properties useful for the calculation of deflection and stresses in beams

In the theory of beams...

... the stiffnesses EA (axial) and $EI = EI_x$ (flexural) are required: the term A represents the area of the cross-section and $I = I_x$ the moment of inertia with respect to a principal centroidal axis (named I_ξ in the following)



CENTROID, FIRST AND SECOND MOMENTS OF AREA





FIRST MOMENTS OF AREA AND CENTROID G

The first moments of area S_x and S_y are calculated as: $A = \int_A dA$ $S_x = \int_A y dA$ $S_y = \int_A x dA$

while the centroid G as:

ARAIA

$$x_{G} = \frac{S_{y}}{A} = \frac{\int_{A} x dA}{A}$$
$$y_{G} = \frac{S_{x}}{A} = \frac{\int_{A} y dA}{A}$$

Environmental Risk Assessment and Mitigation on Cultural Heritage Assets in Central Asia



4





FIRST MOMENTS OF THE AREA AND CENTROID G





- The centroid is the point of an area to the respect of which both first moments of area are null.
- The first moments of the area are null for any pair of centroidal axes
- The first moments may be positive, negative or null; the area A is always positive



SECOND MOMENTS OF THE AREA

The second moments of the area or moments of inertia are

$$I_{xx} = \int_{A} y^{2} dA$$
$$I_{yy} = \int_{A} x^{2} dA$$
$$I_{xy} = \int_{A} xy dA$$



- Ixx and Iyy are always greater than zero
- Ixy (mixed moment of inertia) can be positive, negative or null



PARALLEL-AXIS THEOREMS





PARALLEL-AXIS THEOREMS

The moment of inertia of an area with respect to a given axis is given by:

$$I_{XX} = I_{X_G X_G} + A y_0^2$$
$$I_{YY} = I_{Y_G Y_G} + A x_0^2$$
$$I_{XY} = I_{X_G Y_G} + A x_0 y_0$$

(Formulae by Huygens)



The moments of inertia $I_{x_Gx_G}$ and $I_{y_Gy_G}$...

... are the smallest within all the possible pairs of axes parallels to the original

ones





PRINCIPAL AXES





PRINCIPAL AXES

ARA

ERAMCA



It is possible to find a pair of axes into which $I_{\xi\eta} = 0$ (principal axes), that are rotated of about ϑ (positive if anti-clockwise) with respect to the centroidal axes





• Orientation of the principal axes (ξ , η)

$$\vartheta = \frac{1}{2} \arctan \frac{2 \, I_{x_G y_G}}{I_{y_G y_G} - I_{x_G x_G}}$$

• Moments of inertia with respect to the principal axes (I_{ξ}, I_{η}) :

$$I_{\xi}, I_{\eta} = \frac{1}{2}(I_{X_G X_G} + I_{Y_G Y_G}) \pm \frac{1}{2}\sqrt{(I_{Y_G Y_G} - I_{X_G X_G})^2 + 4 I_{X_G Y_G}^2}$$

choosing I_{ξ} , I_{η} in such a way that:

•
$$I_{\xi} > I_{\eta}$$
 if $I_{x_G x_G} > I_{y_G y_G}$

•
$$I_{\xi} < I_{\eta}$$
 if $I_{x_G x_G} < I_{y_G y_G}$





• If
$$I_{x_G x_G} = I_{y_G y_G}$$
 it is:

$$artheta=+rac{\pi}{4}$$
 and $I_{\xi}=I_{x_Gx_G}-I_{x_Gy_G}$, $I_{\eta}=I_{x_Gx_G}+I_{x_Gy_G}$

Note

- In the principal axes $I_{\xi\eta}=0$
- I_{ξ} and I_{η} are always greater than zero
- Axes both centroidal and principal: central





SIMPLE SECTIONS







$$A = \int_{A} dA = \int_{0}^{b} \int_{0}^{h} dx dy = b h$$
$$S_{x} = \int_{A} y dA = \int_{0}^{b} \int_{0}^{h} y dx dy = +\frac{b h^{2}}{2}$$
$$S_{y} = \int_{A} x dA = \int_{0}^{b} \int_{0}^{h} x dx dy = +\frac{h b^{2}}{2}$$



$$x_G = \frac{S_y}{A} = \frac{+\frac{hb^2}{2}}{bh} = +\frac{b}{2}$$
$$y_G = \frac{S_x}{A} = \frac{+\frac{bh^2}{2}}{bh} = +\frac{h}{2}$$



RECTANGULAR CROSS-SECTION



$$I_{xx} = \int_{A} y^{2} dA = \int_{o}^{b} \int_{o}^{h} y^{2} dx dy = \frac{b h^{3}}{3}$$
$$I_{yy} = \int_{A} x^{2} dA = \int_{o}^{b} \int_{o}^{h} x^{2} dx dy = \frac{h b^{3}}{3}$$
$$I_{xy} = \int_{A} xy dA = \int_{o}^{b} \int_{o}^{h} xy dx dy = +\frac{b^{2} h^{2}}{4}$$

$$I_{x_{G}x_{G}} = I_{xx} - Ay_{G}^{2} = \frac{bh^{3}}{3} - (bh)\left(+\frac{h}{2}\right)^{2} = \frac{bh^{3}}{12}$$
$$I_{y_{G}y_{G}} = I_{yy} - Ax_{G}^{2} = \frac{hb^{3}}{3} - (bh)\left(+\frac{b}{2}\right)^{2} = \frac{hb^{3}}{12}$$
$$I_{x_{G}y_{G}} = I_{xy} - Ax_{G}y_{G} = +\frac{b^{2}h^{2}}{4} - (bh)\left(+\frac{b}{2}\right)\left(+\frac{h}{2}\right) = 0$$

The rectangular cross-section is doubly symmetrical

Co-funded by the ERAMCA of the European Union



COMPOSITE AREAS





PRINCIPAL AXES FOR COMPOSITE AREAS









- Divide the area in *n* simple shapes (which geometrical properties are known, from tables)
- Compute the total area (summing the *n* elementary areas)
- Compute the total first moments (summing the first moments of the *n* elementary areas, keeping in mind the signs) with respect to an arbitrary set of axes (*x*, *y*)
- Determine the position of the centroid



- Compute the total moments of inertia (summing the moments of inertia of the *n* elementary areas) remembering that:
 - the parallel-axis theorem has to be used in order to refer all the moments of inertia to the global centroidal axes (x_G, y_G) ; in tables the m.o.i. are related to the local centroids of the elementary areas (x_{G_i}, y_{G_i})
 - the mixed inertia moment of the *i*-th area depends on the direction of the axes. It is thus necessary to control the directions of the local axes x_{G_i}, y_{G_i} with respect to the global ones x_G, y_G
- Compute the orientation of the principal axes (ϑ) and the moments of inertia related to the principal axes (I_ξ e I_η)



PARTICULAR SYMMETRIES



- Axial symmetry: the centroid is on the symmetry axis and one of the principal axes coincides with the symmetry axis. The other one is perpendicular
- Polar symmetry: the centroid G coincides with the center of the area C



BE CAREFUL...



• Some simple shapes may result misleading...



EXAMPLES





SOLVED EXAMPLE



18

SOLVED EXAMPLE

$$x_{G} = \frac{S_{Y}}{A} = \frac{\sum_{i=1}^{n} S_{Y}^{i}}{\sum_{i=1}^{n} A^{i}} = \frac{\sum_{i=1}^{n} A^{i} x_{G_{i}}}{\sum_{i=1}^{n} A^{i}} = \frac{10.430a^{3}}{6.715a^{2}} = +1.553a$$
$$y_{G} = \frac{S_{X}}{A} = \frac{\sum_{i=1}^{n} S_{X}^{i}}{\sum_{i=1}^{n} A^{i}} = \frac{\sum_{i=1}^{n} A^{i} y_{G_{i}}}{\sum_{i=1}^{n} A^{i}} = \frac{8.323a^{3}}{6.715a^{2}} = +1,239a$$

$$\begin{split} I_{x_G x_G} &= 2.034a^4 + 2.085a^4 = 4.119a^4 \\ I_{y_G y_G} &= 5.201a^4 + 0.160a^4 = 5.361a^4 \\ I_{x_G y_G} &= 0.125a^4 + 0.718a^4 = +0.843a^4 \end{split}$$

$$\vartheta = \frac{1}{2} \arctan \frac{2 I_{x_G y_G}}{I_{y_G y_G} - I_{x_G x_G}} = \frac{1}{2} \arctan \frac{2 \times (+0.843a^4)}{5.361a^4 - 4.119a^4} = +26.8^{\circ} \quad \text{(counterclockwise)}$$
$$I_{\xi}, I_{\eta} = \frac{1}{2} (I_{x_G x_G} + I_{y_G y_G}) \pm \frac{1}{2} \sqrt{(I_{y_G y_G} - I_{x_G x_G})^2 + 4 I_{x_G y_G}^2} = \dots = 4.740a^4 \pm 1.047a^4$$
so that
$$\begin{cases} I_{\xi} = 3.693a^4 \\ I_{\eta} = 5.787a^4 \end{cases}$$



Parameter	Physical dimension	SI unit
x _G , y _G	L	m
А	L ²	m²
S _x , S _y	L ³	m ³
I _{xx} , I _{yy} , I _{xy}	L ⁴	m ⁴
$I_{x_Gx_G}, I_{y_Gy_G}, I_{x_Gy_G}, I_{\xi}, I_{\eta}$	L ⁴	m ⁴

